

# 西崎業績

## 著書

(欧文)

1. Nishizaki T, Kanno T. Resveratrol: A Candidate Drug for Treating Rheumatoid Arthritis. In Rheumatoid Arthritis - Treatment, ISBN 978-953-307-850-2 (Ed. Andrew B), In Tech, 269-284 (2012).
2. Nishizaki T, Ikeuchi Y, Mori M, Okada Y. Adenosine activates the K<sup>+</sup> channel and cytosolic Ca<sup>2+</sup> release via a P2Y purinoceptor in hippocampal neurons. In "The Role of Adenosine in the Nervous System" (Ed. Okada Y), Elsevier, 143-150 (1997).

(和文)

1. 東 英穂, 吉村 恵, 西崎 知之, 福田 淳, 片山 芳文, 佐久間 康夫. CBT 対応病態で学ぶ生理学. 丸善株式会社 (2004).
2. 西崎知之. 脳神経科学イラストレイテッド 第1版. カラーアトラス: 電位・電流記録の基本. 第4章(5) シナプス可塑性. 羊土社 (2000).
3. 西崎知之. 脳神経科学イラストレイテッド 改訂第2版. カラーアトラス: 電位・電流記録の基本. 第8章(1) さまざまな実験法. 羊土社 (2006).

## 総説

(和文)

1. 西崎 知之. 不飽和脂肪酸誘導体 DCP-LA によるシナプス前終末 $\alpha 7$  アセチルコリン受容体を標的とした海馬シナプス伝達促進作用. 生体の科学 2009;60(3):248-55.
2. 岡田 安弘, 浅野 富子, 西崎 知之, 荒川 俊雄. GABA 研究の進歩. 神経精神薬理 1988;10:117-38.

## 原著臨床論文

(欧文)

1. Nishizaki T, Tamaki N, Fujiwara H, Matsumoto S. Posterior fossa subdural effusion due to head trauma. Neurosurg 1988;23(1):81-4.
2. Nishizaki T, Tamaki N, Takeda N, Shirakuni T, Kondoh T, Matsumoto S. Dolichoectatic basilar artery: a review of 23 cases. Stroke 1986;17(6):1277-81.
3. Nishizaki T, Tamaki N, Matsumoto S, Fujita S. Consideration of the operative indications for posterior fossa venous angiomas. Surg Neurol 1986;25:441-5.
4. Nishizaki T, Wakabayashi T. A self-retaining multipore suction tube. J Neurosurg 1985;63:304.
5. Nishizaki T, Tamaki N, Matsumoto S, Fujita S. Multiple giant fusiform aneurysms with hydrocephalus. Surg Neurol 1985;24:101-4.
6. Nishizaki T, Tamaki N, Nishida Y, Matsumoto S. Bilateral internuclear ophthalmoplegia due to hydrocephalus: a case report. Neurosurg 1985;17(5):822-5.
7. Nishizaki T, Tamaki N, Nishida Y, Fujita K, Matsumoto S. Aneurysms of the distal posterior inferior cerebellar artery: experience with three cases and review of the literature. Neurosurg 1985;16(6): 829-32.

(和文)

1. 藤田 勝三, 山下 晴央, 増村 道雄, 西崎 知之, 玉木 紀彦, 松本 悟. Natural history of giant intracranial aneurysms. 脳神経外科 1988;16: 225-31.
2. 西崎 知之, 玉木 紀彦, 白国 隆行, 細田 弘吉, 松本 悟, 藤田 稠清. Giant serpentine aneurysm の 2 例. 臨床放射線 1986;31:309-12.
3. 玉木 紀彦, 白国 隆行, 西崎 知之, 西田 吉充, 松本 悟. 中脳水道閉塞による水頭症. 中外医薬 1984;37:46-50.

## 原著研究論文

(欧文)

1. Nishizaki T, Gotoh Akihiro, Gotoh Akinobu. GSK-3 $\beta$  may be a factor for restraint stress-induced depression-related behaviors and spatial learning impairment independent of tau phosphorylation. *Int J Curr Res* 2022;14(6):21606-21610.
2. Nishizaki T, Tanaka A. PKC $\epsilon$  stimulates vesicular exocytosis of  $\alpha 7$  nicotinic ACh receptor by targeting NSF. *Int J Curr Res* 2019;11(6):4995-5000.
3. Nishizaki T, Mabuchi M, Tanaka A. The anti-dementia drug DCP-LA binds to NSF specifically. *Int J Curr Res* 2019;11(6):4594-4597.
4. Nishizaki T. HUHS1015 reduces PKM2 due to autophagic degradation in gastric cancer cells. *Int J Curr Res* 2019;11(5):3935-3940.
5. Nishizaki T. HUHS1015 downregulates PKM2 through an AMPK/p38MAPK pathway in MKN45 cells. *Int J Curr Res* 2019;11(1):35-39.
6. Nishizaki T. Fe<sup>3+</sup> facilitates endocytic internalization of extracellular A $\beta$ <sub>1-42</sub> and enhances A $\beta$ <sub>1-42</sub>-induced caspase-3/caspase-4 activation and neuronal cell death. *Mol Neurobiol* 2019;56(7):4812-4819.
7. Nishizaki T. PKM2 is a target of HUHS1015 for inducing colonic cancer cell death. *Int J Curr Res* 2018;10(12):76406-76412.
8. Nishizaki T. IL-33 suppresses GSK-3 $\beta$  activation through an ST2-independent MyD88/TRAF6/RIP/PI3K/Akt pathway. *Heliyon* 2018;4(11):e00971.
9. Nishizaki T. Profilin facilitates PKC $\epsilon$  activation by accelerating ATP supply. *Biochem Biophys Res Commun* 2018;506(4):918-922.
10. Nishizaki T. Profilin enhances the activated PKC $\epsilon$  activity and stimulates translocation of  $\alpha 7$  ACh receptor towards the cell surface. *Int J Curr Res* 2018;10(09):73379-73381.
11. Nishizaki T. Nefiracetam and uridine enhance the activated PKC $\epsilon$  activity possibly by lowering the kinase activation energy. *Int J Curr Res* 2018;10(08):72438-72441.
12. Nishizaki T. Enhancement of PKC $\epsilon$  activity through serine autophosphorylation. *Int J Curr Res* 2018;10(5):69726-69730.
13. Nishizaki T. PKC $\epsilon$  activator DCP-LA facilitates assembly of NSF/myosin Va/ $\alpha 7$  ACh receptor. *Int J Curr Res* 2018;10(5):69406-69409.
14. Nishizaki T. Dioleoylphosphoethanolamine retains cell surface GLUT4 by inhibiting PKC $\alpha$ -driven internalization. *Cell Physiol Biochem* 2018;46(5):1985-1998

15. Nishizaki T. PKC $\epsilon$  increases extracellular elastin and fibulin-5/DANCE in dermal fibroblasts. *Cell Physiol Biochem* 2018;46:291-302.
16. Nishizaki T. Linoleic acid derivative DCP-LA prevents Tau phosphorylation by targeting GSK-3 $\beta$ . *Int J Curr Res* 2018;10(01):6425-64430.
17. Nishizaki T. New indomethacin actions linked to cognitive functions. *Int J Curr Res* 2017;9(11):61161-61164.
18. Nishizaki T. Salivary NGF may become a merkmal for early diagnosis of senile dementia. *Ind J Med Res Pharmaceut Sci* 2017;4(1):22-27.
19. Nishizaki T. DCP-LA, a new strategy for Alzheimer's disease therapy. *J Neurol Neuromed* 2017;2(9):1-8.
20. Nishizaki T. Dilinoleoylphosphatidylethanolamine may restrain the progress of MCI into Alzheimer's disease. *Gerontol Geriatr Res* 2017;1(1):113.
21. Nishizaki T. IL-33 acts to express Schaffer collateral/CA1 LTP and regulate learning and memory by targeting MyD88. *Neural Plast* 2017;2017:2531453. doi: 10.1155/2017/2531453.
22. Nishizaki T. Linoleic acid derivative DCP-LA sheds light on treatment of Alzheimer's disease. *Gerontol Geriatr: Res* 2017;3(2):1032.
23. Nishizaki T. Does IL-33 potentiate or depress AMPA receptor response? *Ind J Med Res Pharmaceut Sci* 2017;4(9):42-47.
24. Nishizaki T. Akt activation and GSK-3 $\beta$  inactivation in the hippocampus are accelerated by deleting Il-33. *Int J Biol Pharm Res* 2017;8(4):159-162.
25. Nishizaki T. The linoleic acid derivative DCP-LA sheds light on type 1 DM therapy. *Ind J Med Res Pharmaceut Sci* 2016;3(16):34-40.
26. Nishizaki T. The phosphatidylethanolamine derivative diDCP-LA-PE stimulates vesicular GLUT4 exocytosis by interacting with NSF. *Ind J Med Res Pharmaceut Sci* 2016;3(12):8-14.
27. Nishizaki T. HUHS1015 promotes autophagic XIAP degradation to induce apoptosis of gastrointestinal cancer cells. *Ind J Med Res Pharmaceut Sci* 2016;8(3):8-17.
28. Nishizaki T. DCP-LA exerts an antiaging action on the skin. *Skin Pharmacol Physiol* 2016;29(3):157-160.
29. Nishizaki T, Gotoh A, Shimizu T, Tanaka A. The phosphatidylethanolamine derivative diDCP-LA-PE mimics intracellular insulin signaling. *Sci Rep* 2016;6:27267.
30. Nishizaki T. DCP-LA revives UV irradiation-induced elastic fiber degradation and epithelial hyperplasia. *Int J Biol Pharm Res* 2016;7(3):150-153.
31. Nishizaki T. *N*-Ethylmaleimide dissociates  $\alpha 7$  ACh receptor from a complex with NSF and

- promotes its delivery to the presynaptic membrane. *Neurochem Res* 2016;41(8):2043-2048.
32. Kaku Y, Tsuchiya A, Shimizu T, Tanaka A, Nishizaki T. HUHS1015 suppresses colonic cancer growth by inducing necrosis and apoptosis in association with mitochondrial damage. *Anticancer Res* 2016;36(1):39-48.
  33. Kanno T, Tsuchiya A, Tanaka A, Nishizaki T. Combination of PKC $\epsilon$  activation and PTP1B inhibition effectively suppresses A $\beta$ -induced GSK-3 $\beta$  activation and Tau phosphorylation. *Mol Neurobiol* 2016;53(7):4787-4797.
  34. Jin Y, Tsuchiya A, Kanno T, Nishizaki T. Amyloid- $\beta$  peptide increases cell surface localization of  $\alpha 7$  ACh receptor to protect neurons from amyloid  $\beta$ -induced damage. *Biochem Biophys Res Commun* 2015;468(1-2):157-160.
  35. Ohyama M, Tsuchiya A, Kaku Y, Kanno K, Shimizu T, Tanaka A, Nishizaki T. Phosphatidylinositol derivatives induce gastric cancer cell apoptosis by accumulating AIF and AMID in the nucleus. *Anticancer Res* 2015;35(12):6563-6571.
  36. Tsuchiya A, Kaku Y, Shimizu T, Tanaka A, Nishizaki T. XIAP is a target for HUHS1015-induced caspase-3 activation in CW2 colonic cancer cells. *Int J Biol Pharm Res* 2015;6(12):978-981.
  37. Tsuchiya A, Kaku Y, Nakano T, Nishizaki T. Diarachidonoylphosphoethanolamine induces apoptosis of malignant pleural mesothelioma cells through a Trx/ASK1/p38 MAPK pathway. *J Pharmacol Sci* 2015;129:160-168.
  38. Yamazaki H, Jin Y, Tsuchiya A, Kanno T, Nishizaki T. Adipose-derived stem cell-conditioned medium ameliorates antidepressant-related behaviors in the mouse model of Alzheimer's disease. *Neurosci Lett* 2015;609:53-57.
  39. Ohyama M, Kaku Y, Tsuchiya A, Kanno K, Shimizu T, Tanaka A, Nishizaki T. HUHS1015 induces apoptosis of MKN45 gastric cancer cells by activating caspase-3 in association with mitochondrial damage. *Int J Biol Pharm Res* 2015; 6(10):771-775.
  40. Kanno T, Tsuchiya A, Shimizu T, Mabuchi M, Tanaka A, Nishizaki T. DCP-LA activates cytosolic PKC $\epsilon$  by interacting with the phosphatidylserine binding/associating sites Arg50 and Ile89 in the C2-like domain. *Cell Physiol Biochem* 2015;37(1):193-200.
  41. Tsuchiya A, Nishizaki T. Anticancer effect of adenosine on gastric cancer through diverse signaling pathways. *World J Gastroenterol* 2015;21(39):10931-10935.
  42. Tsuchiya A, Kanno T, Shimizu T, Tanaka A, Nishizaki T. Phosphatidylinositol derivative enantiomers have the potential for the antitumor effect against malignant pleural mesothelioma cells. *Int J Biol Pharm Res* 2015;6(9):731-733

43. Tsuchiya A, Kanno T, Shimizu T, Tanaka A, Nishizaki T. Rac1 and ROCK are implicated in the cell surface delivery of GLUT4 under the control of the insulin signal mimetic diDCP-LA-PE. *J Pharmacol Sci* 2015;128(4):179-184.
44. Tsudaka S, Kanno T, Nishizaki T. Levetiracetam enhances  $\alpha 7$  ACh receptor responses by indirectly interacting with protein kinase A and protein kinase C. *Int J Biol Pharm Res* 2015;6(7):558-562.
45. Kanemura S, Tsuchiya A, Kanno T, Nakano T, Nishizaki T. Phosphatidylinositol induces caspase-independent apoptosis of malignant pleural mesothelioma cells by accumulating AIF in the nucleus. *Cell Physiol Biochem* 2015;36(3):1037-1048
46. Kaku Y, Tsuchiya A, Kanno T, Nakano T, Nishizaki T. Diarachidonoylphosphoethanolamine induces necrosis/necroptosis of malignant pleural mesothelioma cells. *Cell Signal* 2015;27(9):1713-1719.
47. Kaku Y, Tsuchiya A, Kanno T, Nishizaki T. Irinotecan induces cell cycle arrest, but not apoptosis or necrosis, in Caco-2 and CW2 colorectal cancer cell lines. *Pharmacology* 2015;95(3-4):154-159.
48. Kaku Y, Tsuchiya A, Kanno T, Nakao S, Shimizu T, Tanaka A, Nishizaki T. The newly synthesized anticancer drug HUHS1015 is useful for treatment of human gastric cancer. *Cancer Chemother Pharmacol* 2015;75(3):527-535.
49. Nakajima Y, Kanno T, Nagaya T, Kuribayashi K, Nakano T, Gotoh A, Nishizaki T. Adenosine deaminase inhibitor EHNA exhibits a potent anticancer effect against malignant pleural mesothelioma. *Cell Physiol Biochem* 2015;35(1):51-60.
50. Kaku Y, Tsuchiya A, Kanno T, Nishizaki T. HUHS1015 induces necroptosis and caspase-independent apoptosis of MKN28 human gastric cancer cells in association with AMID accumulation in the nucleus. *Anticancer Agents Med Chem* 2015;15(2):242-247.
51. Miyake T, Ito T, Yanai A, Inoue N, Miyagawa Y, Murase K, Imamura M, Ichii S, Takatsuka Y, Nishizaki T, Hirota S, Ohtsuka M, Yamamoto H, Noguchi S, Miyoshi Y. C4.4A highly expressed in HER2-positive human breast cancers may indicate a good prognosis. *Breast Cancer* 2015;22(4):366-373.
52. Nishizaki T, Kanno T, Tsuchiya A, Kaku Y, Shimizu T, Tanaka A. 1-[2-(2-Methoxyphenylamino)ethylamino]-3-(naphthalene-1-yloxy)propan-2-ol may be a promising anticancer drug. *Molecules* 2014;19(12):21462-21472.
53. Tsuchiya A, Nagaya H, Kanno T, Nishizaki T. Oleic acid stimulates glucose uptake into adipocytes by enhancing insulin receptor signaling. *J Pharmacol Sci* 2014;126(4):337-343.

54. Jin Y, Kanno T, Nishizaki T. Acute restraint stress impairs induction of long-term potentiation by activating GSK-3 $\beta$ . *Neurochem Res* 2015;40(1):36-40.
55. Mikami K, Nagaya H, Gotoh A, Kanno A, Tsuchiya A, Nakano T, Nishizaki T. Naftopidil is useful for treatment of malignant pleural mesothelioma. *Pharmacology* 2014;94(3-4):163-169.
56. Kanno T, Tsuchiya A, Nishizaki T. Hyperphosphorylation of Tau at Ser396 occurs in the much earlier stage than appearance of learning and memory disorders in 5XFAD mice. *Behav Brain Res* 2014;274:302-306.
57. Tsuchiya A, Kanno T, Nishizaki T. Dipalmitoleoylphosphoethanolamine as a PP2A enhancer obstructs insulin signaling by promoting Ser/Thr dephosphorylation of Akt. *Cell Physiol Biochem* 2014;34(3):617-627.
58. Kanno T, Jin Y, Nishizaki T. DL-/PO-phosphatidylcholine restores restraint stress-induced depression-related behaviors and spatial memory impairment. *Behav Pharmacol* 2014;25(5-6):575-581.
59. Tian N, Kanno T, Jin Y, Nishizaki T. Lithium potentiates GSK-3 $\beta$  activity by inhibiting phosphoinositide 3-kinase-mediated Akt phosphorylation. *Biochem Biophys Res Commun* 2014;450(1):746-749.
60. Kanno T, Tanaka A, Nishizaki T. Linoleic acid derivative DCP-LA ameliorates stress-induced depression-related behavior by promoting cell surface 5-HT<sub>1A</sub> receptor translocation, stimulating serotonin release, and inactivating GSK-3 $\beta$ . *Mol Neurobiol* 2015;51(2):523-532.
61. Kaku Y, Nagaya H, Tsuchiya A, Kanno T, Gotoh A, Tanaka A, Shimizu T, Tabata C, Nakano T, Nishizaki T. Newly synthesized anticancer drug HUHS1015 is effective on malignant pleural mesothelioma. *Cancer Sci* 2014;105(7):883-889.
62. Tsuchiya A, Kanno T, Nagaya H, Shimizu T, Tanaka A, Nishizaki T. PTP1B inhibition causes Rac1 activation by enhancing receptor tyrosine kinase signaling. *Cell Physiol Biochem* 2014;33(4):1097-1105.
63. Kaku Y, Tsuchiya A, Kanno T, Nakano T, Nishizaki T. Dipalmitoleoyl-phosphatidylethanolamine induces apoptosis of NCI-H28 malignant mesothelioma cells. *Anticancer Res* 2014;34(4):1759-1764.
64. Tsuchiya A, Kanno T, Nishizaki T. Anticancer action of adenosine through diverse signaling pathways. *Personalized Med Univ* 2014;3:35-37.
65. Nishizaki T, Kanno T, Gotoh A. Evaluation of the newly synthesized linoleic acid derivative DCP-LA as a promising anti-dementia drug. *Personalized Med Univ* 2014;3:28-34.

66. Kanno T, Gotoh A, Nishizaki T. Sphingosine induces apoptosis and cell cycle arrest by targeting SDK and PKC $\delta$ . *Personalized Med Univ* 2014;3:22-27.
67. Tsuchiya A, Kanno T, Shimizu T, Nakao S, Tanaka A, Nishizaki T. DCP-LA-phosphatidylinositol and its enantiomer exhibit different bioactivities. *Cell Physiol Biochem* 2014;33(2):300-309.
68. Tsuchiya A, Kanno T, Shimizu T, Nakao S, Tanaka A, Tabata C, Nakano T, Nishizaki T. A novel PP2A enhancer induces caspase-independent apoptosis of MKN28 gastric cancer cells with high MEK activity. *Cancer Lett* 2014;347(1):123-128.
69. Murase K, Yanai A, Saito M, Imamura M, Miyagawa Y, Takatsuka Y, Inoue N, Ito T, Hirota S, Sasa M, Katagiri T, Fujimoto Y, Hatada T, Ichii S, Nishizaki T, Tomita N, Miyoshi Y. Biological characteristics of luminal subtypes in pre- and postmenopausal estrogen receptor-positive and HER2-negative breast cancers. *Breast Cancer* 2014;21(1):52-57.
70. Tsuchiya A, Kanno T, Nishizaki T. Stearic acid serves as a potent inhibitor of protein tyrosine phosphatase 1B. *Cell Physiol Biochem* 2013;32(5):1451-1459.
71. Tsuchiya A, Kanno T, Nishizaki T. PI3 kinase directly phosphorylates Akt1/2 at Ser473/474 in the insulin signal transduction pathway. *J Endocrinol* 2013;220(1):49-59.
72. Shibata E, Kanno T, Tsuchiya A, Kuribayashi K, Tabata C, Nakano T, Nishizaki T. Free fatty acids inhibit protein tyrosine phosphatase 1B and activate Akt. *Cell Physiol Biochem* 2013;32(4):871-879.
73. Mikami K, Tabata C, Tabata R, Nogi Y, Terada T, Honda M, Kamiya H, Nishizaki T, Nakano T. Clinical significance of serum angiopoietin-1 in malignant peritoneal mesothelioma. *Cancer Invest* 2013;31(8):511-515.
74. Niba ET, Nagaya H, Kanno T, Tsuchiya A, Gotoh A, Tabata C, Kuribayashi K, Nakano T, Nishizaki T. Crosstalk between PI3 kinase/PDK1/Akt/Rac1 and Ras/Raf/MEK/ERK pathways downstream PDGF receptor. *Cell Physiol Biochem* 2013;31(6):905-913.
75. Kanno T, Tanaka A, Shimizu T, Nakano T, Nishizaki T. 1-[2-(2-Methoxyphenylamino)ethylamino]-3-(naphthalene-1-yloxy)propan-2-ol as a potential anticancer drug. *Pharmacology* 2013;91(5-6):339-345.
76. Tsuchiya A, Kanno T, Nishizaki T. Diacylglycerol promotes GLUT4 translocation to the cell surface in a PKC $\epsilon$ -dependent and PKC $\lambda/\iota$  and  $-\zeta$ -independent manner. *Life Sci* 2013;93(5-6):240-246.
77. Tabata C, Kanemura S, Tabata R, Masachika E, Shibata E, Otsuki T, Nishizaki T, Nakano T. Serum HMGB1 as a diagnostic marker for malignant peritoneal mesothelioma. *J Clin Gastroenterol* 2013;47(8):684-688.



78. Tabata C, Shibata E, Tabata R, Kanemura S, Mikami K, Nogi Y, Masachika E, Nishizaki T, Nakano T. Serum HMGB1 as a prognostic marker for malignant pleural mesothelioma. *BMC Cancer* 2013;13:205.
79. Kanno T, Tsuchiya A, Shimizu T, Nakao S, Tanaka A, Nishizaki T. Effects of newly synthesized DCP-LA-phospholipids on protein kinase C and protein phosphatases. *Cell Physiol Biochem* 2013;31(4-5):555-564.
80. Masachika E, Kanno T, Nakano T, Gotoh A, Nishizaki T. Naftopidil induces apoptosis in malignant mesothelioma cell lines independently of  $\alpha_1$ -adrenoceptor blocking. *Anticancer Res* 2013;33(3):887-894.
81. Gotoh A, Nagaya H, Kanno T, Tagawa M, Nishizaki T. Fiber-substituted conditionally replicating adenovirus Ad5F35 induces oncolysis of human bladder cancer cells in in vitro analysis. *Urology* 2013;81(4):920.e7-11.
82. Kanno T, Tsuchiya A, Tanaka A, Nishizaki T. The linoleic acid derivative DCP-LA increases membrane surface localization of the  $\alpha_7$  ACh receptor in a protein 4.1N-dependent manner. *Biochem J* 2013;450(2):303-309.
83. Nishizaki T, Kanno T, Gotoh A. DL-/PO-phosphatidylcholine may shed bright light on treatment of Alzheimer dementia. *Personalized Med Univ* 2013;2:12-15.
84. Nagaya H, Gotoh A, Kanno T, Nishizaki T. A<sub>3</sub> adenosine receptor mediates apoptosis in in vitro RCC4-VHL human renal cancer cells by up-regulating AMID expression. *J Urol* 2013;189(1):321-328.
85. Nishimoto T, Kanno T, Shimizu T, Tanaka A, Nishizaki T. Regulation of GluA1 AMPA receptor through PKC phosphorylation induced by free fatty acid derivative HUHS2002. *Lipids* 2013;48(1):23-28.
86. Okuwa H, Kanno T, Fujita Y, Gotoh A, Tabata C, Fukuoka K, Nakano T, Nishizaki T. Sphingosine suppresses mesothelioma cell proliferation by inhibiting PKC- $\delta$  and inducing cell cycle arrest at the G<sub>0</sub>/G<sub>1</sub> phase. *Cell Physiol Biochem* 2012;30(4):995-1004.
87. Kanno T, Nishimoto T, Fujita Y, Gotoh A, Nakano T, Nishizaki T. Sphingosine induces apoptosis in MKN-28 human gastric cancer cells in an SDK-dependent manner. *Cell Physiol Biochem* 2012;30(4):987-994.
88. Kanno T, Tsuchiya A, Shimizu T, Tanaka A, Nishizaki T. Indomethacin serves as a potential inhibitor of protein phosphatases. *Cell Physiol Biochem* 2012;30(4):1014-1022.

89. Kanno T, Gotoh A, Fujita Y, Nakano T, Nishizaki T. A<sub>3</sub> adenosine receptor mediates apoptosis in 5637 human bladder cancer cells by G<sub>q</sub> protein/PKC-dependent AIF upregulation. *Cell Physiol Biochem* 2012;30(5):1159-1168.
90. Kanno T, Gotoh A, Nakano T, Tagawa M, Nishizaki T. Beneficial oncolytic effect of fiber-substituted conditionally replicating adenovirus on human lung cancer. *Anticancer Res* 2012;32(11):4891-4895.
91. Kanno T, Nishizaki T. A<sub>2a</sub> adenosine receptor mediates PKA-dependent glutamate release from synaptic-like vesicles and Ca<sup>2+</sup> efflux from an IP<sub>3</sub>- and ryanodine-insensitive intracellular calcium store in astrocytes. *Cell Physiol Biochem* 2012;30(6):1398-1412.
92. Gotoh A, Nagaya H, Kanno T, Nishizaki T. Antitumor action of α<sub>1</sub>-adrenoceptor blockers on human bladder, prostate and renal cancer cells. *Pharmacology* 2012;90(5-6):242-246.
93. Gotoh A, Kanno T, Nagaya H, Nakano T, Tabata C, Fukuoka K, Tagawa M, Nishizaki T. Gene therapy using adenovirus against malignant mesothelioma. *Anticancer Res* 2012;32(9):3743-3747.
94. Kanno T, Nakano T, Fujita Y, Gotoh A, Nishizaki T. Adenosine induces apoptosis in SBC-3 human lung cancer cells through A<sub>3</sub> adenosine receptor-dependent AMID upregulation. *Cell Physiol Biochem* 2012;30(3):666-677.
95. Kanno T, Tsuchiya A, Shimizu T, Tanaka A, Nishizaki T. Novel indomethacin action: selective and direct activation of protein kinase C-ε. *Cell Physiol Biochem*. 2012;30(3):771-777.
96. Kanno T, Shimizu T, Tanaka A, Nishimoto T, Nishizaki T. Free fatty acid derivative HUHS2002 potentiates α<sub>7</sub> ACh receptor responses through indirect activation of CaMKII. *Lipids* 2012;47(9):865-871.
97. Otsuki T, Kanno T, Fujita Y, Tabata C, Fukuoka K, Nakano T, Gotoh A, Nishizaki T. A<sub>3</sub> adenosine receptor-mediated p53-dependent apoptosis in Lu-65 human lung cancer cells. *Cell Physiol Biochem* 2012;30(1):210-220.
98. Kanno T, Tanaka A, Nishizaki T. Linoleic acid derivative DCP-LA stimulates vesicular transport of α<sub>7</sub> ACh receptors towards surface membrane. *Cell Physiol Biochem* 2012;30(1):75-82.
99. Nogi Y, Kanno T, Nakano T, Fujita Y, Tabata C, Fukuoka K, Gotoh A, Nishizaki T. AMP converted from intracellularly transported adenosine upregulates p53 expression to induce malignant pleural mesothelioma cell apoptosis. *Cell Physiol Biochem* 2012;30(1):61-74.
100. Nagaya H, Tagawa M, Hiwasa K, Terao S, Kanno T, Nishizaki T, Gotoh A. Fiber-substituted conditionally replicating adenovirus for oncolysis of human renal carcinoma cells. *Anticancer Res* 2012;32(7):2985-2989.

101. Kamiya H, Kanno T, Fujita Y, Gotoh A, Nakano T, Nishizaki T. Apoptosis-related gene transcription in human A549 lung cancer cells via A<sub>3</sub> adenosine receptor. *Cell Physiol Biochem* 2012;29(5-6):687-696.
102. Honda M, Kanno T, Fujita Y, Gotoh A, Nakano T, Nishizaki T. Mesothelioma cell proliferation through autocrine activation of PDGF- $\beta\beta$  receptor. *Cell Physiol Biochem* 2012;29(5-6):667-674.
103. Kanno T, Yaguchi T, Shimizu T, Tanaka A, Nishizaki T. 8-[2-(2-Pentyl-cyclopropylmethyl)-cyclopropyl]-octanoic acid and its diastereomers improve age-related cognitive deterioration. *Lipids* 2012;47(7):687-695.
104. Okada A, Yaguchi T, Kanno T, Gotoh A, Nakano T, Nishizaki T. PDGF-D/PDGF- $\beta\beta$  receptor-regulated chemotaxis of malignant mesothelioma cells. *Cell Physiol Biochem* 2012;29(1-2):241-250.
105. Kanno T, Yaguchi T, Nagata T, Nishizaki T. Indomethacin activates protein kinase C and potentiates  $\alpha 7$  ACh receptor responses. *Cell Physiol Biochem* 2012;29(1-2):189-196.
106. Tsuchiya A, Kanno T, Saito M, Miyoshi Y, Gotoh A, Nakano T, Nishizaki T. Intracellularly transported adenosine induces apoptosis in MCF-7 human breast cancer cells by accumulating AMID in the nucleus. *Cancer Lett* 2012;321(1):65-72.
107. Tamura K, Kanno T, Fujita Y, Gotoh A, Nakano T, Nishizaki T. A<sub>2a</sub> adenosine receptor mediates HepG2 cell apoptosis by downregulating Bcl-X<sub>L</sub> expression and upregulating Bid expression. *J Cell Biochem.* 2012;113(5):1766-1775.
108. Kanno T, Yaguchi T, Nagata T, Shimizu T, Tanaka A, Nishizaki T. Indomethacin enhances learning and memory potential by interacting with CaMKII. *J Cell Physiol* 2012;227(3):919-926.
109. Nakayama H, Yaguchi T, Yoshiya S, Nishizaki T. Resveratrol induces apoptosis MH7A human rheumatoid arthritis synovial cells in a sirtuin 1-dependent manner. *Rheumatol Int* 2012;32(1):151-157.
110. Yaguchi T, Nakano T, Gotoh A, Nishizaki T. Adenosine promotes GATA-2-regulated p53 gene transcription to induce HepG2 cell apoptosis. *Cell Physiol Biochem* 2011;28(4):761-770.
111. Kanno T, Nishizaki T. CFTR mediates noradrenaline-induced ATP efflux from DRG neurons. *Mol Pain* 2011;7:72.
112. Kanno T, Nishizaki T. Endogenous sphingosine 1-phosphate regulates spontaneous glutamate release from mossy fiber terminals via S1P<sub>3</sub> receptors. *Life Sci* 2011;89(3-4):137-140.
113. Kanno T, Nishizaki T. Sphingosine induces apoptosis in hippocampal neurons and astrocytes by activating caspase-3/-9 via a mitochondrial pathway linked to SDK/14-3-3 protein/Bax/cytochrome c. *J Cell Physiol* 2011;226(9):2329-2337.

114. Shimizu T, Kanno T, Tanaka A, Nishizaki T.  $\alpha,\beta$ -DCP-LA selectively activates PKC- $\epsilon$  and stimulates neurotransmitter release with the highest potency among 4 diastereomers. *Cell Physiol Biochem* 2011;27(2):149-158.
115. Yang D, Yaguchi T, Nakano T, Nishizaki T. Adenosine activates AMPK to phosphorylate Bcl-X<sub>L</sub> responsible for mitochondrial damage and DIABLO release in HuH-7 cells. *Cell Physiol Biochem* 2011;27(1):71-78.
116. Yang D, Yaguchi T, Nagata T, Gotoh A, Dovat S, Song C, Nishizaki T. AMID mediates adenosine-induced caspase-independent HuH-7 cell apoptosis. *Cell Physiol Biochem* 2011;27(1):37-44.
117. Nagata T, Yaguchi T, Nishizaki T. DL- and PO-phosphatidylcholines as a promising learning and memory enhancer. *Lipids Health Dis* 2011;10:25.
118. Suzuki T, Murakami K, Izuo N, Kume T, Akaike A, Nagata T, Nishizaki T, Tomiyama T, Mori H, Irie K. E22 $\Delta$  mutation in amyloid  $\beta$ -protein promotes  $\beta$ -sheet transformation, radical production, and synaptotoxicity, but not neurotoxicity. *Int J Alzheimers Dis* 2011;doi:10.4061/2011/431320.
119. Mabuchi M, Haramura M, Shimizu T, Nishizaki T, Tanaka A. Selective elution of target protein from affinity resins by a simple reductant with a thiol group. *Bioorg Med Chem Lett* 2010;20(24):7361-7364.
120. Kanno T, Nishizaki T, Proia RL, Kajimoto T, Jahangeer S, Okada T, Nakamura S. Regulation of synaptic strength by sphingosine 1-phosphate in the hippocampus. *Neuroscience* 2010;171(4):973-980.
121. Yaguchi T, Nagata T, Nishizaki T. 1,2-Dilinoleoyl-sn-glycero-3- phosphoethanolamine ameliorates age-related spatial memory deterioration by preventing neuronal cell death. *Behav Brain Funct* 2010;6:52.
122. Yamada K, Yaguchi T, Kanno T, Mukasa T, Nishizaki T. Auto-positive feedback regulation for nicotinic acetylcholine receptors by protein kinase C activation. *Cell Physiol Biochem* 2010;26(2):247-252.
123. Yaguchi T, Saito M, Yasuda Y, Kanno T, Nakano T, Nishizaki T. Higher concentrations of extracellular ATP suppress proliferation of Caco-2 human colonic cancer cells via an unknown receptor involving PKC inhibition. *Cell Physiol Biochem* 2010;26(2):125-134.
124. Yaguchi T, Nishizaki T. Extracellular high K<sup>+</sup> stimulates vesicular glutamate release from astrocytes by activating voltage-dependent calcium channels. *J Cell Physiol* 2010;225(2):512-518.
125. Kanno T, Yaguchi T, Nishizaki T. Noradrenaline stimulates ATP release from DRG neurons by targeting  $\beta_3$  adrenoceptors as a factor of neuropathic pain. *J Cell Physiol* 2010;224(2):345-351.

126. Kanno T, Yaguchi T, Nagata T, Mukasa T, Nishizaki T. Regulation of AMPA receptor trafficking by *O*-glycosylation. *Neurochem Res* 2010;35(5):782-788.
127. Yaguchi T, Fujikawa H, Nishizaki T. Linoleic acid derivative DCP-LA protects neurons from oxidative stress-induced apoptosis by inhibiting caspase-3/-9 activation. *Neurochem Res* 2010;35(5):712-717.
128. Yang D, Yaguchi T, Nakano T, Nishizaki T. Adenosine-induced caspase-3 activation by tuning Bcl-X<sub>L</sub>/DIABLO/IAP expression in HuH-7 human hepatoma cells. *Cell Biol Toxicol* 2010;26(4):319-330.
129. Yaguchi T, Saito M, Yasuda Y, Nishizaki T. Caspase-4 activation in association with decreased adenosine deaminase activity may be a factor for gastric ulcer. *Digestion* 2010;81(1):62-67.
130. Yang D, Yaguchi T, Lim CR, Ishizawa Y, Nakano T, Nishizaki T. Tuning of apoptosis-mediator gene transcription in HepG2 human hepatoma cells through an adenosine signal. *Cancer Lett* 2010;291(2):225-229.
131. Saito M, Yaguchi T, Yasuda Y, Nakano T, Nishizaki T. Adenosine suppresses CW2 human colonic cancer growth by inducing apoptosis via A<sub>1</sub> adenosine receptors. *Cancer Lett* 2010;290(2):211-215.
132. Yaguchi T, Nagata T, Yang D, Nishizaki T. Interleukin-18 regulates motor activity, anxiety and spatial learning without affecting synaptic plasticity. *Behav Brain Res* 2010;206(1):47-51.
133. Nagata T, Tomiyama T, Mori H, Yaguchi T, Nishizaki T. DCP-LA neutralizes mutant amyloid  $\beta$  peptide-induced impairment of long-term potentiation and spatial learning. *Behav Brain Res* 2010;206(1):151-154.
134. Yaguchi T, Muramoto M, Nakano T, Nishizaki T. Urinary trypsin inhibitor suppresses migration of malignant mesothelioma. *Cancer Lett* 2010;288(2):214-218.
135. Kanno T, Yaguchi T, Nagata T, Tanaka A, Nishizaki T. DCP-LA stimulates AMPA receptor exocytosis through CaMKII activation due to PP-1 inhibition. *J Cell Physiol* 2009;221(1):183-188.
136. Yaguchi T, Nagata T, Nishizaki T. 1-Palmitoyl-2-oleoyl-*sn*-glycero- 3-phosphocholine improves cognitive decline by enhancing long-term depression. *Behav Brain Res* 2009;204(1):129-132.
137. Nakagomi T, Taguchi A, Fujimori Y, Saino O, Nakano-Doi A, Kubo S, Gotoh A, Soma T, Yoshikawa H, Nishizaki T, Nakagomi N, Stern DM, Matsuyama T. Isolation and characterization of neural stem/progenitor cells from post-stroke cerebral cortex in mice. *Eur J Neurosci* 2009;29(9):1842-1852.
138. Yamamoto S, Kanno T, Yamada K, Yasuda Y, Nishizaki T. Dual regulation of heat-activated K<sup>+</sup> channel in rat DRG neurons via  $\alpha_1$  and  $\beta$  adrenergic receptors. *Life Sci.* 2009;85(3-4):167-171.

139. Yasuda Y, Saito M, Yamamura T, Yaguchi T, Nishizaki T. Extracellular adenosine induces apoptosis in Caco-2 human colonic cancer cells by activating caspase-9/-3 via A<sub>2a</sub> adenosine receptors. *J Gastroenterol* 2009;44(1):56-65.
140. Yaguchi T, Nagata T, Nishizaki T. Dilinoleoylphosphatidylethanolamine ameliorates scopolamine-induced impairment of spatial learning and memory by targeting  $\alpha 7$  nicotinic ACh receptors. *Life Sci* 2009;84(9-10):263-266.
141. Fujikawa H, Kanno T, Nagata T, Nishizaki T. The phosphodiesterase III inhibitor olprinone inhibits hippocampal glutamate release via a cGMP/PKG pathway. *Neurosci Lett* 2008;448(2):208-211.
142. Watanabe K, Kanno T, Oshima T, Miwa H, Tashiro C, Nishizaki T. Vagotomy upregulates expression of the *N*-methyl-D-aspartate receptor NR2D subunit in the stomach. *J Gastroenterol* 2008;43(5):322-326.
143. Watanabe K, Kanno T, Oshima T, Miwa H, Tashiro C, Nishizaki T. The NMDA receptor NR2A subunit regulates proliferation of MKN45 human gastric cancer cells. *Biochem Biophys Res Commun* 2008;367(2):487-490.
144. Tomiyama T, Nagata T, Shimada H, Teraoka R, Fukushima A, Kanemitsu H, Takuma H, Kuwano R, Imagawa M, Ataka S, Wada Y, Yoshioka E, Nishizaki T, Watanabe Y, Mori H. A new amyloid  $\beta$  variant favoring oligomerization in Alzheimer's-type dementia. *Ann Neurol* 2008;63(3):377-387.
145. Yang D, Yaguchi T, Yamamoto H, Nishizaki T. Intracellularly transported adenosine induces apoptosis in HuH-7 human hepatoma cells by downregulating c-FLIP expression causing caspase-3/-8 activation. *Biochem Pharmacol* 2007;73(10):1665-1675.
146. Sakamoto T, Ebisu Y, Ikeda A, Nakaya M, Nishizaki T. Hippocampal size may contribute prospective diagnosis of age-related dementia. *Psychogeriatrics* 2007;7(2):76-80.
147. Watanabe K, Yaguchi T, Yang D, Kanno T, Nagai K, Yamamoto S, Fujikawa H, Yamamoto H, Nagata T, Tashiro C, Nishizaki T. Beneficial effect of intracellular free high-mannose oligosaccharides on cryopreservation of mammalian cells and proteins. *Cryobiology* 2006;53(3):330-335.
148. Kanno T, Yamamoto H, Yaguchi T, Hi R, Mukasa T, Fujikawa H, Nagata T, Yamamoto S, Tanaka A, Nishizaki T. The linoleic acid derivative DCP-LA selectively activates PKC- $\epsilon$ , possibly binding to the phosphatidylserine binding site. *J Lipid Res* 2006;47(6):1146-1156.
149. Sai K, Yang D, Yamamoto H, Fujikawa H, Yamamoto S, Nagata T, Saito M, Yamamura T, Nishizaki T. A<sub>1</sub> adenosine receptor signal and AMPK involving caspase-9/-3 activation are

- responsible for adenosine-induced RCR-1 astrocytoma cell death. *Neurotoxicology* 2006;27(4):458-467.
150. Maruo K, Yamamoto S, Kanno T, Yaguchi T, Maruo S, Yashiya S, Nishizaki T. Tunicamycin decreases the probability of single-channel openings for *N*-methyl-D-aspartate and  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid receptors. *Neuroreport* 2006;17(3):313-317.
  151. Yaguchi T, Nagata T, Mukasa T, Fujikawa H, Yamamoto H, Yamamoto S, Iso H, Tanaka A, Nishizaki T. Linoleic acid derivative DCP-LA improves learning impairment in SAMP8. *Neuroreport* 2006;17(1):105-108.
  152. Maruo K, Yamamoto H, Yamamoto S, Nagata T, Fujikawa H, Kanno T, Yaguchi T, Maruo S, Yoshiya S, Nishizaki T. Modulation of P2X receptors via adrenergic pathways in rat dorsal root ganglion neurons after sciatic nerve injury. *Pain* 2006;120(1-2):106-112.
  153. Kanno T, Yaguchi T, Yamamoto S, Nagata T, Yamamoto H, Fujikawa H, Nishizaki T. Bidirectional regulations for glutamate and GABA release in the hippocampus by  $\alpha 7$  and non- $\alpha 7$  ACh receptors. *Biochem Biophys Res Commun* 2005;338(2):742-747.
  154. Kanno T, Yaguchi T, Yamamoto S, Yamamoto H, Fujikawa H, Nagata T, Tanaka A, Nishizaki T. 8-[2-(2-Pentyl-cyclopropylmethyl)-cyclopropyl]-octanoic acid stimulates GABA release from interneurons projecting to CA1 pyramidal neurons in the rat hippocampus via pre-synaptic  $\alpha 7$  acetylcholine receptors. *J Neurochem* 2005;95(3):695-702.
  155. Yaguchi T, Yamamoto S, Nagata T, Kanno T, Tanaka A, Nishizaki T. Effects of *cis*-unsaturated free fatty acids on PKC- $\epsilon$  activation and nicotinic ACh receptor responses. *Mol Brain Res* 2005;133(2):320-324.
  156. Nagata T, Yamamoto S, Yaguchi T, Iso H, Tanaka A, Nishizaki T. The newly synthesized linoleic acid derivative DCP-LA ameliorates memory deficits in animal models treated with amyloid- $\beta$  peptide and scopolamine. *Psychogeriatrics* 2005;5(4):122-126.
  157. Yamamoto S, Kanno T, Nagata T, Yaguchi T, Tanaka A, Nishizaki T. The linoleic acid derivative FR236924 facilitates hippocampal synaptic transmission by enhancing activity of presynaptic  $\alpha 7$  acetylcholine receptors on the glutamatergic terminals. *Neuroscience* 2005;130(1):207-213.
  158. Kanno T, Nagata T, Yamamoto S, Okamura H, Nishizaki T. Interleukin-18 stimulates synaptically released glutamate and enhances postsynaptic AMPA receptor responses in the CA1 region of mouse hippocampal slices. *Brain Res* 2004;1012(1-2):190-193.

159. Saitoh M, Nagai K, Nakagawa K, Yamamura T, Yamamoto S, Nishizaki T. Adenosine induces apoptosis in the human gastric cancer cells via an intrinsic pathway relevant to activation of AMP-activated protein kinase. *Biochem Pharmacol* 2004;67(10):2005-2011.
160. Nishizaki T. ATP- and adenosine-mediated signaling in the central nervous system: adenosine stimulates glutamate release from astrocytes via A<sub>2a</sub> adenosine receptors. *J Pharmacol Sci* 2004;94(2):100-102.
161. Ohta K, Miyamoto H, Yaguchi T, Nagai K, Yamamoto S, Nomura T, Tanaka A, Nishizaki T. Stearic acid facilitates hippocampal neurotransmission by enhancing nicotinic ACh receptor responses via a PKC pathway. *Mol Brain Res* 2003;119(1):83-89.
162. Saitoh M, Nagai K, Yaguchi T, Fujikawa Y, Ikejiri K, Yamamoto S, Nakagawa K, Yamamura T, Nishizaki T. Arachidonic acid peroxides induce apoptotic Neuro-2A cell death in association with intracellular Ca<sup>2+</sup> rise and mitochondrial damage independently of caspase-3 activation. *Brain Res* 2003;991(1-2):187-194.
163. Miyamoto H, Yaguchi T, Ohta K, Nagai K, Nagata T, Yamamoto S, Nishizaki T. 2-Pyrrolidinone induces a long-lasting facilitation of hippocampal synaptic transmission by enhancing  $\alpha 7$  ACh receptor responses via a PKC pathway. *Mol Brain Res* 2003;117(1):91-96.
164. Matsumoto S, Yamamoto S, Sai K, Maruo K, Adachi M, Saitoh M, Nishizaki T. Pipelicolic acid induces apoptosis in neuronal cells. *Brain Res* 2003;980(2):179-184.
165. Maruo K, Nagata T, Yamamoto S, Nagai K, Yajima Y, Maruo S, Nishizaki T. Tunicamycin inhibits NMDA and AMPA receptor responses independently of *N*-glycosylation. *Brain Res* 2003;977(2):294-297.
166. Nishizaki T. *N*-Glycosylation sites on the nicotinic ACh receptor subunits regulate receptor channel desensitization and conductance. *Mol Brain Res* 2003;114(2):172-176.
167. Tanaka A, Nishizaki T. The newly synthesized linoleic acid derivative FR236924 induces a long-lasting facilitation of hippocampal neurotransmission by targeting nicotinic acetylcholine receptors. *Bioorg Med Chem Lett* 2003;13(6):1037-1040.
168. Tada H, Uchino M, Nagai K, Nomura T, Kondoh T, Saito N, Yamamura T, Yajima Y, Nishizaki T. The anti-dementia drug FK960 stimulates glial glutamate release via a PKA pathway. *Mol Brain Res* 2002;109(1-2):63-68.
169. Nagai K, Jiang MH, Hada J, Nagata T, Yajima Y, Yamamoto S, Nishizaki T. (-)-Epigallocatechin gallate protects against NO stress-induced neuronal damage after ischemia by acting as an anti-oxidant. *Brain Res* 2002;956(2):319-322.



170. Matsumoto S, Kanno T, Nomura T, Shiotani T, Watabe S, Nabeshima T, Nishizaki T. Nefiracetam induces a long-lasting facilitation of hippocampal postsynaptic responses in mice lacking the NMDA receptor  $\epsilon 1$  subunit. *Brain Res* 2002;956(1):178-181.
171. Aihara H, Fujiwara S, Mizuta I, Tada H, Kanno T, Tozaki H, Nagai K, Yajima Y, Inoue K, Kondoh T, Motooka Y, Nishizaki T. Adenosine triphosphate accelerates recovery from hypoxic/hypoglycemic perturbation of guinea pig hippocampal neurotransmission via a  $P_2$  receptor. *Brain Res* 2002;952(1):31-37.
172. Ohta K, Nomura T, Kanno T, Nagai K, Yamamoto S, Yajima Y, Kondoh T, Kohmura E, Saito N, Nishizaki T. L-trans-PDC enhances hippocampal neuronal activity by stimulating glial glutamate release independently of blocking transporters. *Biochem Biophys Res Commun* 2002;295(2):376-381.
173. Nishizaki T, Nagai K, Nomura T, Tada H, Kanno T, Tozaki H, Li XX, Kondoh T, Kodama N, Takahashi E, Sakai N, Tanaka K, Saito N. A new neuromodulatory pathway with a glial contribution mediated via  $A_{2a}$  adenosine receptors. *Glia* 2002;39(2):133-147.
174. Tozaki H, Matsumoto A, Kanno T, Nagai K, Nagata T, Yamamoto S, Nishizaki T. The inhibitory and facilitatory actions of amyloid- $\beta$  peptides on nicotinic ACh receptors and AMPA receptors. *Biochem Biophys Res Commun* 2002;294(1):42-45.
175. Nishizaki T, Matsumura T. The aniracetam metabolite 2-pyrrolidinone induces a long-term enhancement in AMPA receptor responses via a CaMKII pathway. *Mol Brain Res* 2002;98(1-2):130-134.
176. Tozaki H, Kanno T, Nomura T, Kondoh T, Kodama N, Saito N, Aihara H, Nagata T, Matsumoto S, Ohta K, Nagai K, Yajima Y, Nishizaki T. Role of glial glutamate transporters in the facilitatory action of FK960 on hippocampal neurotransmission. *Mol Brain Res*. 2001;97(1):7-12.
177. Aihara H, Li X, Fujiwara S, Matsumura T, Okumura S, Tozaki H, Kanno T, Ohta K, Nagai K, Nishizaki T. The protective action of nefiracetam against electrophysiological and metabolic damage in the hippocampus after deprivation of glucose and oxygen. *Brain Res* 2001;922(1):158-162.
178. Motooka Y, Kondoh T, Nomura T, Tamaki N, Tozaki H, Kanno T, Nishizaki T. Selective cholinergic denervation inhibits expression of long-term potentiation in the adult but not infant rat hippocampus. *Dev Brain Res* 2001;129(1):119-123.
179. Nomura T, Nishizaki T, Enomoto T, Itoh H. A long-lasting facilitation of hippocampal neurotransmission via a phospholipase  $A_2$  signaling pathway. *Life Sci* 2001;68(25):2885-2891.

180. Li XX, Nomura T, Aihara H, Nishizaki T. Adenosine enhances glial glutamate efflux via A<sub>2a</sub> adenosine receptors. *Life Sci* 2001;68(12):1343-1350.
181. Kondoh T, Nishizaki T, Aihara H, Tamaki N. NMDA-responsible, APV-insensitive receptor in cultured human astrocytes. *Life Sci* 2001;68(15):1761-1767.
182. Nishizaki T, Nomura T, Matsuyama S, Kondoh T, Fujimoto E, Yoshii M. Critical role of nicotinic ACh receptor in the formation of long-term potentiation: Implications for development of anti-dementia drugs. *Psychogeriatrics* 2001;1(3):209-217.
183. Yoshii M, Watabe S, Murashima YL, Nukada T, Shiotani T, Nishizaki T. Neuronal Ca<sup>2+</sup> channels and nicotinic ACh receptors as functional target of the nootropic nefiracetam. *Psychogeriatrics* 2001;1(1):39-49.
184. Nishizaki T, Nomura T, Matuoka T, Kondoh T, Enikolopov G, Sumikawa K, Watabe S, Shiotani T, Yoshii M. The anti-dementia drug nefiracetam facilitates hippocampal synaptic transmission by functionally targeting presynaptic nicotinic ACh receptors. *Mol Brain Res* 2000;80(1):53-62.
185. Matsuyama S, Matsumoto A, Enomoto T, Nishizaki T. Activation of nicotinic acetylcholine receptors induces long-term potentiation *in vivo* in the intact mouse dentate gyrus. *Eur J Neurosci* 2000;12(10):3741-3747.
186. Nomura T, Nishizaki T. Nefiracetam facilitates hippocampal neurotransmission by a mechanism independent of the piracetam and aniracetam action. *Brain Res* 2000;870(1-2):157-162.
187. Nishizaki T, Matsuoka T, Nomura T, Kondoh T, Watabe S, Shiotani T, Yoshii M. Presynaptic nicotinic acetylcholine receptors as a functional target of nefiracetam in inducing a long-lasting facilitation of hippocampal neurotransmission. *Alzheimer Dis Assoc Disord* 2000;14 Suppl 1:S82-94.
188. Matsuyama S, Nomura T, Nishizaki T. Sustained facilitatory action of FK960 on hippocampal neurotransmission. *Brain Res* 2000;857(1-2):317-320.
189. Ishihara I, Minami Y, Nishizaki T, Matsuoka T, Yamamura H. Activation of calpain precedes morphological alterations during hydrogen peroxide-induced apoptosis in neuronally differentiated mouse embryonal carcinoma P19 cell line. *Neurosci Lett* 2000;279(2):97-100.
190. Shimada F, Shiga Y, Morikawa M, Kawazura H, Morikawa O, Matsuoka T, Nishizaki T, Saito N. The neuroprotective agent MS-153 stimulates glutamate uptake. *Eur J Pharmacol* 1999;386(2-3):263-270.
191. Nishizaki T, Matsuoka T, Sumikawa K. Positive- and negative-regulation of Ca<sup>2+</sup> entry through ACh receptor channels by phosphorylation. *Mol Brain Res* 1999;69(2):295-301.

192. Nishizaki T, Nomura T, Matsuoka T, Enikolopov G, Sumikawa K. Arachidonic acid induces a long-lasting facilitation of hippocampal synaptic transmission by modulating PKC activity and nicotinic ACh receptors. *Mol Brain Res* 1999;69(2):263-272.
193. Nishizaki T, Matsuoka T, Nomura T, Kondoh T, Tamaki N, Okada Y. Store Ca<sup>2+</sup> depletion enhances NMDA responses in cultured human astrocytes. *Biochem Biophys Res Commun* 1999;259(3):661-664.
194. Nishizaki T, Matsuoka T, Nomura T, Matsuyama S, Watabe S, Shiotani T, Yoshii M. A 'long-term-potential-like' facilitation of hippocampal synaptic transmission induced by the nootropic nefiracetam. *Brain Res* 1999;826(2):281-288.
195. Nishizaki T, Matsuoka T, Nomura T, Enikolopov G, Sumikawa K. Arachidonic acid potentiates currents through Ca<sup>2+</sup>-permeable AMPA receptors by interacting with a CaMKII pathway. *Mol Brain Res* 1999;67(1):184-189.
196. Nishizaki T, Nomura T, Matsuoka T, Tsujishita Y. Arachidonic acid as a messenger for the expression of long-term potentiation. *Biochem Biophys Res Commun* 1999;254(2):446-449.
197. Nishizaki T, Sumikawa K. Nicotinic receptors are regulated by protein kinase C activated via a nicotinic receptors-mediated signaling pathway. *Mol Brain Res* 1998;61(1-2):211-218.
198. Matsuoka T, Nishizaki T, Nomura T, Mori M, Okada Y. ATP produces potassium currents via P<sub>3</sub> purinoceptor in the follicle cell layer of *Xenopus* oocytes. *Neurosci Lett* 1998;248(2):130-132.
199. Nishizaki T, Matsuoka T, Nomura T, Sumikawa K. Modulation of ACh receptor currents by arachidonic acid. *Mol Brain Res* 1998;57(1):173-179.
200. Nishizaki T, Mori M. Diverse signal transduction pathways mediated by endogenous P<sub>2</sub> receptors in cultured rat cerebral cortical neurons. *J Neurophysiol* 1998;79(5):2513-2521.
201. Nishizaki T, Matsuoka T. Low glucose enhances Na<sup>+</sup>/glucose transport in bovine brain artery endothelial cells. *Stroke* 1998;29(4):844-849.
202. Matsuoka T, Nishizaki T, Kisby GE. Na<sup>+</sup>-dependent and phlorizin-inhibitable transport of glucose and cycasin in brain endothelial cells. *J Neurochem* 1998;70(2):772-777.
203. Nishizaki T, Matsuoka T, Nomura T, Sumikawa K, Shiotani T, Watabe S, Yoshii M. Nefiracetam modulates acetylcholine receptor currents via two different signal transduction pathways. *Mol Pharmacol* 1998;53(1):1-5.
204. Nomura T, Nishizaki T. Methamphetamine modulates ACh-evoked currents in *Xenopus* oocytes expressing the rat  $\alpha 7$  receptors. *Neurosci Lett* 1997;239(2-3):73-76.
205. Nishizaki T, Sumikawa K. Lysophosphatidic acid potentiates ACh receptor currents by G-protein-mediated activation of protein kinase C. *Mol Brain Res* 1997;50(1-2):121-126.

206. Matsuoka T, Nishizaki T, Nomura T. The voltage-dependent non-selective cation channel sensitive to the L-type calcium channel blocker efonidipine regulates Ca<sup>2+</sup> influx in brain vascular smooth muscle cells. *Biochem Biophys Res Commun* 1997;240(2):484-487.
207. Nishizaki T, Matsuoka T, Nomura T, Sumikawa K. A serum factor potentiates ACh and AMPA receptor currents via differential signal transduction pathways. *Biochem Biophys Res Commun* 1997;238(2):565-568.
208. Matsuoka T, Kondoh T, Tamaki N, Nishizaki T. The GABA<sub>A</sub> receptor is expressed in human neurons derived from a teratocarcinoma cell line. *Biochem Biophys Res Commun* 1997;237(3):719-723.
209. Mori M, Hosomi H, Nishizaki T, Kawahara K, Okada Y. Calcium release from intracellular stores evoked by extracellular ATP in a *Xenopus* renal epithelial cell line. *J Physiol* 1997;502 ( Pt 2):365-373.
210. Ikeuchi Y, Nishizaki T, Matsuoka T, Sumikawa K. Long-lasting enhancement of ACh receptor currents by lysophospholipids. *Mol Brain Res* 1997;45(2):317-320.
211. Nishizaki T, Ikeuchi Y, Matsuoka T, Sumikawa K. Short-term depression and long-term enhancement of ACh-gated channel currents induced by linoleic and linolenic acid. *Brain Res* 1997;751(2):253-258.
212. Matsuoka T, Nishizaki T, Ikeuchi Y, Okada Y, Sumino K. Differential effects of the serotonin receptors on cultured rat cerebral cortical neurons. *Life Sci* 1997;53(3):233-236.
213. Matsuoka T, Nishizaki T, Sumino K. A specific inhibitory action of lithium on the 5-HT<sub>2c</sub> receptor expressed in *Xenopus laevis* oocytes. *Mol Pharmacol* 1997;51(3):471-474.
214. Nishizaki T, Ikeuchi Y, Matsuoka T, Sumikawa K. Oleic acid enhances ACh receptor currents by activation of Ca<sup>2+</sup>/calmodulin-dependent protein kinase II. *Neuroreport* 1997;8(3):597-601.
215. Gehle VM, Walcott EC, Nishizaki T, Sumikawa K. *N*-glycosylation at the conserved sites ensures the expression of properly folded functional ACh receptors. *Mol Brain Res* 1997;45(2):219-229.
216. Ikeuchi Y, Nishizaki T, Mori M, Okada Y. Adenosine activates the K<sup>+</sup> channel and enhances cytosolic Ca<sup>2+</sup> release via a P2Y purinoceptor in hippocampal neurons. *Eur J Pharmacol* 1996;304(1-3):191-199.
217. Ikeuchi Y, Nishizaki T. ATP-regulated K<sup>+</sup> channel and cytosolic Ca<sup>2+</sup> mobilization in cultured rat spinal neurons. *Eur J Pharmacol* 1996;302(1-3):163-169.
218. Ikeuchi Y, Nishizaki T, Matsuoka T, Sumikawa K. Arachidonic acid potentiates ACh receptor currents by protein kinase C activation but not by receptor phosphorylation. *Biochem Biophys Res Commun* 1996;221(3):716-721.

219. Mori M, Nishizaki T, Kawahara K, Okada Y. ATP-activated cation conductance in a *Xenopus* renal epithelial cell line. *J Physiol* 1996;491 ( Pt 2):281-290.
220. Ikeuchi Y, Nishizaki T, Okada Y. Repetitive applications of ATP potentiate potassium current by  $Ca^{2+}$ /calmodulin kinase in cultured rat hippocampal neurons. *Neurosci Lett* 1996;203(2):115-118.
221. Matsuoka T, Nishizaki T, Ikeuchi Y. Regulation of the serum-activated  $Ca^{2+}$ -dependent chloride channel in *Xenopus* oocytes. *Biochem Biophys Res Commun* 1996;218(2):633-637.
222. Ikeuchi Y, Nishizaki T, Mori M, Okada Y. Regulation of the potassium current and cytosolic  $Ca^{2+}$  release induced by 2-methylthio ATP in hippocampal neurons. *Biochem Biophys Res Commun* 1996;218(2):428-433.
223. Ikeuchi Y, Nishizaki T.  $P_2$  purinoceptor-operated potassium channel in rat cerebellar neurons. *Biochem Biophys Res Commun* 1996;218(1):67-71.
224. Nishizaki T, Ikeuchi Y. Adenosine evokes potassium currents by protein kinase C activated via a novel signaling pathway in superior colliculus neurons. *FEBS Lett* 1996;378(1):1-6.
225. Ikeuchi Y, Nishizaki T, Matsuoka T. Lysophosphatidylcholine inhibits NMDA-induced currents by a mechanism independent of phospholipase  $A_2$ -mediated protein kinase C activation in hippocampal glial cells. *Biochem Biophys Res Commun* 1995;217(3):811-816.
226. Ikeuchi Y, Nishizaki T. Dual effects of ATP on the potassium channel and intracellular  $Ca^{2+}$  release in smooth muscle cells of the bovine brain arteries. *Biochem Biophys Res Commun* 1995;215(3):1071-1077.
227. Ikeuchi Y, Nishizaki T. ATP activates the potassium channel and enhances cytosolic  $Ca^{2+}$  release via a  $P_2Y$  purinoceptor linked to pertussis toxin-insensitive G-protein in brain artery endothelial cells. *Biochem Biophys Res Commun* 1995;215(3):1022-1028.
228. Ikeuchi Y, Nishizaki T, Mori M, Okada Y. Adenosine activates the potassium channel via a  $P_2$  purinoceptor but not via an adenosine receptor in cultured rat superior colliculus neurons. *Neurosci Lett* 1995;198(3):205-208.
229. Ikeuchi Y, Nishizaki T, Okada Y. A  $P_2$  purinoceptor activated by ADP in rat medullar neurons. *Neurosci Lett* 1995;198(2):71-74.
230. Ikeuchi Y, Nishizaki T. The  $P_2Y$  purinoceptor-operated potassium channel is possibly regulated by the beta gamma subunits of a pertussis toxin-insensitive G-protein in cultured rat inferior colliculus neurons. *Biochem Biophys Res Commun* 1995;214(2):589-596.
231. Nishizaki T, Ikeuchi Y. Activation of endogenous protein kinase C enhances currents through  $\alpha 1$  and  $\alpha 2$  glycine receptor channels. *Brain Res* 1995;687(1-2):214-216.

232. Ikeuchi Y, Nishizaki T. Methylcobalamin induces a long-lasting enhancement of the postsynaptic field potential in hippocampal slices of the guinea pig. *Neurosci Lett* 1995;192(2):113-116.
233. Ikeuchi Y, Nishizaki T. ATP-evoked potassium currents in rat striatal neurons are mediated by a P<sub>2</sub> purinergic receptor. *Neurosci Lett* 1995;190(2):89-92.
234. Nishizaki T, Kammesheidt A, Sumikawa K, Asada T, Okada Y. A sodium- and energy-dependent glucose transporter with similarities to SGLT1-2 is expressed in bovine cortical vessels. *Neurosci Res* 1995;22(1):13-22.
235. Nishizaki T, Sumikawa K. Direct action of 4- $\beta$ -phorbol-12,13-dibutyrate (PDBu) on nicotinic acetylcholine receptor channel independent of protein kinase C activation. *Neurosci Lett* 1995;188(2):129-131.
236. Sumikawa K, Nishizaki T. The amino acid residues 1-128 in the  $\alpha$  subunit of the nicotinic acetylcholine receptor contain assembly signals. *Mol Brain Res* 1994;25(3-4):257-264.
237. Nishizaki T, Sumikawa K. A cAMP-dependent Ca<sup>2+</sup> signalling pathway at the endplate provided by the  $\gamma$  to  $\epsilon$  subunit switch in ACh receptors. *Mol Brain Res* 1994;24(1-4):341-345.
238. Nishizaki T, Morales A, Gehle VM, Sumikawa K. Differential interactions of gentamicin with mouse junctional and extrajunctional ACh receptors expressed in *Xenopus* oocytes. *Mol Brain Res* 1994;21(1-2):99-106.
239. Nishizaki T, Sumikawa K. Tunicamycin alters channel gating characteristics of junctional and extrajunctional acetylcholine receptors expressed in *Xenopus* oocytes. *Neurosci Lett* 1994;170(2):273-276.
240. Nishizaki T, Sumikawa K. Tunicamycin increases desensitization of junctional and extrajunctional acetylcholine receptors expressed in *Xenopus* oocytes by a mechanism independent of *N*-glycosylation blocking. *Mol Pharmacol* 1992;42(1):152-156.
241. Mori M, Nishizaki T, Okada Y. Protective effect of adenosine on the anoxic damage of hippocampal slice. *Neuroscience* 1992;46(2):301-307.
242. Nishizaki T, Okada Y. Effects of excitatory amino acids on the oxygen consumption of hippocampal slices from the guinea pig. *Brain Res* 1988;452(1-2):11-20.
243. Nishizaki T, Yamauchi R, Tanimoto M, Okada Y. Effects of temperature on the oxygen consumption in thin slices from different brain regions. *Neurosci Lett* 1988;86(3):301-305.
244. Nishizaki T, Yamauchi R, Okada Y. Enhancement of the oxygen consumption in the hippocampal slices of the guinea pig induced by glutamate and its related substances. *Neurosci Lett* 1988;85(1):61-64.